

Preon Trinity

Jean-Jacques Dugne,¹ Sverker Fredriksson,² Johan Hansson,² and Enrico Predazzi³

¹*Laboratoire de Physique Corpusculaire,*

Université Blaise Pascal, Clermont-Ferrand II, FR-63177 Aubière, France

²*Department of Physics, Luleå University of Technology, SE-97187 Luleå, Sweden*

³*Department of Theoretical Physics, Università di Torino, IT-10125 Torino, Italy*

We present a new minimal model for the substructure of all known quarks, leptons and weak gauge bosons, based on only three fundamental and stable spin-1/2 preons. As a consequence, we predict three new quarks, three new leptons, and six new vector bosons. One of the new quarks has charge $-4e/3$. The model explains the apparent conservation of three lepton numbers, as well as the so-called Cabibbo-mixing of the d and s quarks, and predicts electromagnetic decays or oscillations between the neutrinos $\bar{\nu}_\mu$ (ν_μ) and ν_e ($\bar{\nu}_e$). Other neutrino oscillations, as well as rarer quark mixings and CP violation can come about due to a small quantum-mechanical mixing of two of the preons in the quark and lepton wave functions.

PACS numbers: 12.60.Rc, 13.35.-r, 14.60.-z, 14.65.-q

The arguments in favor of a substructure of quarks and leptons in terms of more fundamental preons have been discussed for two decades (a review is given in [1]). The most compelling ones are that there exist so many quarks and leptons, and that they seemingly form a pattern of three “families”.

Much work in the field has been inspired also by the fact that the weak gauge bosons, Z^0 and $W^\pm$, can be understood as being composite. Hence, the weak force would not be fundamental, but rather a “leakage” of multi-preon states, in analogy to the nuclear force being mediated by quark-antiquark states. If so, there is no fundamental electroweak unification, and no theoretical complications caused by massive, unstable and electrically charged gauge bosons. Consequently, there is no need for the so-called Higgs mechanism with its many by-products, *e.g.*, “multiple Higgs”, “composite Higgs”, and “higgsinos”, motivated mainly by internal theoretical problems [2].

Another hint at a substructure is that most quarks and leptons are *unstable*, which in our view disqualifies them as fundamental particles. Historically, all decays of “elementary” particles have sooner or later been interpreted in terms of processes among even more fundamental objects. This wisdom has not yet been included in the so-called standard model, where all quarks and leptons are considered fundamental, stable or not.

Preon models have so far focused on explaining the lightest family of two quarks (u and d) and two leptons (e and ν_e) with as few preons as possible. This means either a minimal number of (two) different preons, *e.g.*, the “rishons” [3,4], or a minimal number of (two) preons inside a quark or lepton, *e.g.*, the “haplons” [5].

The two heavier families are usually considered “excitations” of the light one, but no simple, consistent and predictive model exists. Neither have preon models (or the standard model) been able to explain the conservation of three different lepton numbers, the mixing of quarks, but not of leptons, as described by the Cabibbo-

Kobayashi-Maskawa formalism [6] (quantified by the so-called CKM matrix), and the CP violation in neutral K meson decays.

A classical problem with all preon models (including ours) is that neutrinos are so much lighter than the quarks, especially as preons are expected to have masses in the TeV range. This is related to a lack of understanding of the forces that keep quarks and leptons together, and also to the problem of defining a mass of a permanently bound object.

The essence of our new preon model is that we explain *all* known quarks and leptons with the help of a minimal number of preons, while maintaining as much symmetry and simplicity as possible. We have been inspired by the haplon model of Fritzsch and Mandelbaum [5], where preons have spin 1/2 and 0. In addition, we guess that there is a pairwise (“supersymmetric”) relation between preons of different spin as far as their charges and composite states are concerned.

It turns out that one needs only three preons of each spin to build all known leptons and quarks. We call the spin-1/2 preons α , β and δ , and the spin-0 ones x , y and z . There are several reasons to believe that the scalar preons (“spreons”) are indeed composite themselves, being spin-0 bound states (“di-preons”) of two spin-1/2 preons in a symmetric way. The notion of a spreon, which refers to names like “squark” and “slepton” within the established theory of supersymmetry (SUSY), is only to remind about a purely phenomenological symmetry in our model. It is similar to the “supersymmetry” between quarks and diquarks, as sometimes quoted in quark-diquark models of baryons [7].

It is worth noting that even if the spreons were truly fundamental, and SUSY partners of preons, the symmetry would *not* be implemented among quarks and leptons. Hence, “squarks” and “sleptons” need not exist, and the real SUSY partners of quarks and leptons would themselves be quarks and leptons!

Our full preon scheme, containing three fundamental

spin-1/2 particles, reads:

charge	$+e/3$	$-2e/3$	$+e/3$
spin-1/2 preons	α	β	δ
spin-0 “spreons”	$x = (\bar{\beta}\bar{\delta})$	$y = (\bar{\alpha}\bar{\delta})$	$z = (\bar{\alpha}\bar{\beta})$

We assume that α , β and δ carry a quantum chromodynamic (QCD) color charge (*anticolor*), *i.e.*, the preon colors are the $\mathbf{3}^*$ representation of $SU(3)_{color}$, when the quark colors are $\mathbf{3}$. By construction, also the x , y and z are then color- $\mathbf{3}^*$ ($\mathbf{3} \otimes \mathbf{3} = \mathbf{3}^* \oplus \mathbf{6}$). With quarks and leptons as, respectively, preon-spreon and preon-antispreon states, the colour representations correctly become $\mathbf{3}$ for quarks and $\mathbf{1}$ for leptons, as they should. We assume that the other mathematically allowed color configurations do not exist in nature.

The preon assignments we propose are summarised in the table below, where the result of combining a preon with a spreon or an antispreon can be found in the respective cells.

	$\bar{x}$ ($\beta\delta$)	$\bar{y}$ ($\alpha\delta$)	$\bar{z}$ ($\alpha\beta$)	x ($\beta\bar{\delta}$)	y ($\bar{\alpha}\bar{\delta}$)	z ($\bar{\alpha}\bar{\beta}$)
α	ν_e	μ^+	ν_τ	u	s	c
β	e^-	$\bar{\nu}_\mu$	τ^-	d	X	b
δ	$\bar{\nu}_{\kappa 1}$	κ^+	$\bar{\nu}_{\kappa 2}$	t	g	h

There are alternative schemes for building the known quarks and leptons, but these turn out to be less consistent in details, and will not be discussed here.

In the remainder of this work, we discuss several aspects of this model, starting with the trivial observation that *all* known quarks and leptons are reproduced with their correct charges, color-charges and spins.

Our scheme does not have the three-family structure, so much quoted, but not understood, within the standard model. There is virtually a pairwise relation between quarks and leptons, but not with the same pairs as conventionally defined. The true mathematical structure should instead be “ $SU(3)_{preon}$ ”, *i.e.*, a preon version of the flavor- $SU(3)$ adopted for the three lightest quarks (u , d and s), although the symmetry must be severely broken, as far as quark and lepton masses are concerned.

As the preons are flavor- $\mathbf{3}$, the three spreons are $\mathbf{3}$ too. Hence, the leptons are $\mathbf{3} \otimes \mathbf{3}^* = \mathbf{1} \oplus \mathbf{8}$, and the quarks are $\mathbf{3} \otimes \mathbf{3} = \mathbf{3}^* \oplus \mathbf{6}$. From the table above it is obvious that the lepton flavor-singlet $\mathbf{1}$ is to be found among the three neutrinos ($\nu_e, \bar{\nu}_\mu, \bar{\nu}_{\kappa 2}$), along the diagonal. It is not possible to pinpoint any particular one, or a linear combination, as the singlet, or make an exact analogy to the baryon octet in the quark model, because of the symmetry breaking. It is less ambiguous that the three diagonal quarks (u, X, h) are flavor- $\mathbf{3}^*$, while the remaining ones are flavor- $\mathbf{6}$.

The scheme contains *three new leptons* and *three new quarks*, which could all be too heavy to be created in current experiments, but naturally of interest for future

accelerator energies in the TeV range. Among both leptons and quarks there are two clear mass-trends. Firstly, the bottom row contains objects that are much heavier than the others, which must be caused by a “naked” superheavy δ preon. The presence of a δ does not, however, make a spreon superheavy, since $(\alpha\beta)$ seems to be heavier than $(\beta\delta)$ and $(\alpha\delta)$. Secondly, there is an increasing mass along the diagonals from upper left to lower right.

The t quark is “predicted” to be superheavy, which explains why it is so much heavier than its conventionally assigned partners, the b quark and the τ lepton. These enormous mass differences are hard to accept as *ad hoc* consequences of the Higgs mechanism in the standard model.

One of the predicted new quarks is the X of charge $-4e/3$, which seemingly does not belong to the superheavy group. This naturally poses a problem to the model, and the apparent lack of experimental indications must be analysed.

First, we note that quarks belonging to different flavor- $SU(3)$ representations need not have related masses. Neither would quark masses need to be simply related to lepton masses. Therefore, it might well happen that the flavor- $\mathbf{3}^*$ quarks have rapidly increasing masses, from u to X and h . Then the quark masses would depend not only on the underlying preon masses, but also strongly on the exact composition of the internal wave functions. Ultimately, all $SU(3)$ representations need not be realised in nature. The rule of the game is, however, that if one object exists, so should the full representation, which makes it difficult to claim that the X “need not exist”.

A second alternative is that the X quark might indeed be the one that is commonly believed to be the top quark, and that the conventional charge assignment ($+2e/3$) of the “top” is incorrect. This heretic idea cannot easily be dismissed, since there is no absolute experimental evidence that the top charge is really $+2e/3$ [8]. The top quark was found through its presumed main decay channels, namely semi-leptonic ones like $t \rightarrow b + \ell^+ + \nu$, or non-leptonic ones like $t \rightarrow b + u + \bar{d}$, where a few b quarks have been “tagged” by a charged muon in semi-leptonic decays. However, the situation is rather complex because a total event contains the decay of the full $t\bar{t}$ pair into several leptons and hadronic jets, and the identification and matching of those are non-trivial. In addition, there are individual events that seemingly contradict the assumption of top-quark production. It can therefore not be excluded that a produced “ b ” is, in fact, an $\bar{b}$ (and *vice versa*), or that a tagged b has been matched with the wrong lepton, *i.e.*, from a W decay of the other quark. These are crucial questions, since the corresponding decay channels for the antiquark $\bar{X}$ would be $\bar{X} \rightarrow \bar{b} + \ell^+ + \nu$ and $\bar{X} \rightarrow \bar{b} + u + \bar{d}$, so that the full $X\bar{X}$ decay would give the same leptons and jets as in a $t\bar{t}$ decay, although with a different $b - W$ matching. This puzzle would require a full new analysis of the “top” data, where all matchings

are remade with the assumption of $X\bar{X}$ decay.

A third, and even more speculative idea, is that a “light” X quark has escaped discovery. All searches for a new quark seem to focus on a “fourth-family” b' with charge $-e/3$ [9]. They also rely on model-dependent assumptions on b' decay, *e.g.*, through a flavor-changing neutral current in $b' \rightarrow b + \gamma$ [10] (searched for with gamma detectors). Such experiments cannot see a decay of the X as given above. Neither has there been systematic searches for new resonances in e^+e^- collisions at high energies in the traditional way of fine-tuning the total energy in small steps. It is, on the other hand, hard to believe that a light X would not have been detected in the search for the top quark [8], unless there is some (narrow) kinematic region that has not been properly covered. Also, an X quark lighter than the W^- should have been seen through decays like $W^- \rightarrow X + \bar{d}$ in the W branching ratios [9]. The case for an X with a mass below, say, the W mass is therefore weak.

Turning now to *lepton decays*, we note that all the known ones can be described as regroupings of preons into less massive states. A decaying charged lepton transfers its spreon to a neutrino, while its preon hides inside a W or Z^0 . As an example, the decay $\mu^- \rightarrow \nu_\mu + e^- + \bar{\nu}_e$ is the preon process $\bar{\alpha}(\bar{\alpha}\bar{\delta}) \rightarrow \bar{\beta}(\bar{\alpha}\bar{\delta}) + \beta(\beta\delta) + \bar{\alpha}(\bar{\beta}\bar{\delta})$, where the latter two leptons are the decay products of $W^- = \beta\bar{\alpha}$ (more on vector bosons below).

The observation of three conserved lepton numbers (L_e, L_μ, L_τ) in all known weak decays is a straightforward consequence of *preon stability*. Therefore, this model is to the best of our knowledge the only one that explains the empirical lepton-number conservation from a first principle. This feature is, however, more or less accidental for those preon regroupings that correspond to the known decays, and there is no general conservation of lepton number in the model. A simple example is the decay of the superheavy κ , *e.g.*, through $\kappa^+ \rightarrow \mu^+ + \nu_e + \bar{\nu}_\tau$, violating all three lepton numbers! Such violations are related to processes that require a break-up of a spreon, and a regrouping of its two preons into other spreons.

Other possible lepton-number violating processes are neutrino *oscillations* and *decays*, if at least one neutrino is massive. Indeed, the two superheavy neutrinos *must* decay, an example being $\bar{\nu}_{\kappa 2} \rightarrow \tau^- + \bar{\nu}_\tau + \mu^+$. Such decays relieve us from problems with cosmic, superheavy and stable neutrinos surviving from the Big Bang, which would have given too much cosmic dark matter.

An interesting observation is that the three neutrinos $\nu_e, \bar{\nu}_\mu$ and $\bar{\nu}_{\kappa 2}$ have *identical* preon content, differing only in the grouping into spreons: $\nu_e = \alpha(\beta\delta)$, $\bar{\nu}_\mu = \beta(\alpha\bar{\delta})$ and $\bar{\nu}_{\kappa 2} = \delta(\alpha\beta)$. This opens up for $\nu_e \leftrightarrow \bar{\nu}_\mu$ oscillations or *electromagnetic* decays like $\bar{\nu}_\mu \rightarrow \nu_e + \gamma$, which would be the equivalent of the decay $\Sigma^0 \rightarrow \Lambda^0 + \gamma$ in the quark model. It should be noted that the unorthodox transition between an *antineutrino* and another neutrino is not forbidden by helicity conservation, since a massive neutrino

has both left-handed and right-handed components, and since the photon carries away helicity. It should also be observed that *charged* leptons cannot decay electromagnetically in this model.

Neutrino oscillations can occur also if there is a quantum-mechanical mixing of the α and δ preons inside spreons, mixing the $(\alpha\beta)$ with the $(\beta\delta)$ in the quark and lepton wave functions. This is not forbidden by any known principles, and would lead to an oscillation between the ν_e and the ν_τ . As will be argued below for quark decays, there are reasons to believe that such a mixing occurs on the *per mille* level. This would also mix the e with the τ by as much, but that would hardly be observable, since the electron is stable. (The best way to observe that two particles are mixtures of the same wave-function components is to measure branching ratios when they decay to one and the same ground state.)

In a future work we will analyse if any of these oscillations or electromagnetic decays are in line with the (not fully consistent) experimental data on neutrino “transitions”, *e.g.*, empirical limitations on decays [11], the suppression of atmospheric muon neutrinos [12], and the recent evidence [13] of a $\nu_\mu \rightarrow \nu_e$ “oscillation”.

Quark decays are known to differ substantially from those of leptons, in the sense that some quarks clearly mix quantum-mechanically with each other when decaying. This behavior can be accounted for, but not explained, by the standard model, in terms of the CKM matrix [6]. In particular, it is not understood why there is only one substantial (“Cabibbo”) mixing, *i.e.*, of d and s , while the others are much smaller.

All this can be qualitatively understood in our model, but probably not in any other model where quarks and leptons are related. The crucial property of quarks is that they are composed of both preons and antipreons, giving access to preon-antipreon *annihilation* channels between certain quarks, or equivalently, giving some quarks identical net preon flavors. In particular, $d = \beta(\bar{\beta}\bar{\delta})$ and $s = \alpha(\bar{\alpha}\bar{\delta})$ have the net flavors of the $\bar{\delta}$ and will mix via $\alpha\bar{\alpha} \leftrightarrow \beta\bar{\beta}$ (mediated by a composite Z^0 , or by any gauge boson). The effect is weak, due to either the high Z^0 mass or a strong di-preon binding, which suppresses the $\alpha\bar{\alpha}$ (and $\beta\bar{\beta}$) wave-function overlaps. There are similar mixing channels between c and t , and between b and g .

The smaller CKM matrix elements cannot be understood in the same way. Instead, a weak $\alpha - \delta$ mixing might be required. It suffices to assume a mixing of only the deeply bound preons inside a di-preon. This will mix u with c , d with b , and t with h .

It is possible that also CP violation can be explained by such a mixing. An interesting observation is that the K^0 and $\bar{K}^0$ mesons have identical preon content, although in different preon/spreon configurations. This is no longer true if we introduce a small $\alpha - \delta$ mixing. If this involves also a quantum-mechanical phase, it should result in a CP violation in the K^0 system. The D^0 and B^0 mesons

do *not* have preon contents that are identical to those of their antiparticles, whatever that means for their possible CP violation.

The *vector bosons* come about as bound preon-antipreon states. The most likely configurations are $W^- = \beta\bar{\alpha}$, $Z^0 = (\alpha\bar{\alpha} - \beta\bar{\beta})/\sqrt{2}$ and $W^+ = \alpha\bar{\beta}$, in analogy with the spin-1 ρ mesons in the quark model. There also exist two heavier and orthogonal combinations (Z' and Z'') of $\alpha\bar{\alpha}$, $\beta\bar{\beta}$ and $\delta\bar{\delta}$ (like the ω and ϕ in the quark model). In addition, there are two neutral states with $\alpha\bar{\delta}$ and $\delta\bar{\alpha}$, as well as two charged states of $\beta\bar{\delta}$ and $\delta\bar{\beta}$ (like the four K^* mesons). There should be mixings among these states, especially as they are all heavy and unstable. One can restrain these mixings by fits to the known W and Z decay modes, but that would be beyond the scope of this first qualitative analysis. It should be added that the lightest Z' could be lighter than the lower (model-dependent) mass limit quoted in [9].

In models with composite W and Z one also expects *scalar* partners. The fact that they have not been observed can be blamed either on high masses or on very weak couplings to quarks and leptons [1]. The latter seems more realistic since spin-0 systems are normally lighter than those with spin 1. The similarity with the mesons in the quark model, and the considerable mass difference between the π and ρ mesons, make us suspect that the missing scalars are rather light, and hence interesting to search for in existing experimental data. One possibility is that some of the heavier scalar mesons are indeed hybrids of quark-antiquark and preon-antipreon states.

In conclusion, our model provides qualitative explanations of several phenomena that are not even addressed in the standard model, or in other composite models. Most of them have to do with the (electro)weak interaction, which is the most troublesome part of the standard model, with its many *ad hoc* parameters and hypothetical particles and processes.

However, many problems remain to be solved before it can be considered also a quantitative alternative. One is the mass-ordering of quarks and leptons, which does not follow the flavor- $SU(3)$ quark-model recipe of the light baryons. Obviously, the masses are determined not only by those of the deeply bound preons, but also of the exact structure of the composed objects. It seems as if the “net” quantum numbers are important, so that the colored quarks are heavier than the color-neutral leptons, and the normal neutrinos are light because they carry almost no quantum properties. The $\bar{\nu}_{\kappa 1}$ and $\bar{\nu}_{\kappa 2}$ are different due to the very heavy, “naked” δ .

There is also a long way to go before we understand how preons bind into quarks and leptons, which is related to the mass problem. Clearly, the strong binding into di-preons must be due to attractive spin forces, *e.g.*, color-magnetism in terms of normal QCD or some even more fundamental interaction. The binding of a preon

with a di-preon into quarks is harder to understand. It is not possible to introduce a new hyper-color, and require that quarks be “hyper-confined” in hyper-color singlets. Rather it seems as if this binding is due to a weaker (hyper)color-electric force, which could be attractive in both color-singlet (lepton) and color-triplet (quark) configurations. Then the quarks would be bound but unconfined, and, in this sense, less “fundamental” than leptons.

Finally, it would be interesting to speculate why nature has provided two preons (α and δ) that differ only in mass. Is there some other property that separates them, and are there more superheavy preons to be “discovered”, *e.g.*, a heavier version also of the β ?

Still, as long as preon models have not yet developed into something as complicated as the standard model (with the Higgs mechanism), we consider them worthy of continued theoretical and experimental efforts. Many issues, such as the existence of heavy quarks, leptons and vector bosons, will hopefully be clarified by the next generation of high-energy accelerators.

We acknowledge an illuminating correspondence with H. Fritzsch, as well as valuable experimental information from R. Partridge and G. VanDalen. This project is supported by the European Commission under contract CHRX-CT94-0450, within the network “The Fundamental Structure of Matter”.

-
- [1] I.A. D’Souza, C.S. Kalman, *Preons* (Singapore, World Scientific 1992).
 - [2] For a review of the Higgs mechanism, see B.A. Kniehl, Phys. Rep. **240**, 211 (1994).
 - [3] H. Harari, Phys. Lett. **86B**, 83 (1979); H. Harari, N. Seiberg, Phys. Lett. **98B**, 269 (1981); Nucl. Phys. B **204**, 141 (1982).
 - [4] M.A. Shupe, Phys. Lett. **86B**, 87 (1979).
 - [5] H. Fritzsch, G. Mandelbaum, Phys. Lett. **102B**, 319 (1981).
 - [6] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi, T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
 - [7] For a review of diquarks, see M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson, D.B. Lichtenberg, Rev. Mod. Phys. **65**, 1199 (1993).
 - [8] F. Abe *et al.*, Phys. Rev. Lett. **74**, 2626 (1995); S. Abachi *et al.*, Phys. Rev. Lett. **74**, 2632 (1995).
 - [9] R.M. Barnett *et al.*, *Review of Particle Physics*, Phys. Rev. D **54**, 1 (1996).
 - [10] S. Abachi *et al.*, Phys. Rev. Lett. **78**, 3818 (1997); and references therein for other searches of new quarks.
 - [11] For a review of empirical information on neutrino decay, see S. Sarkar, to be published in Proc. XVI Int. Workshop on Weak Interactions and Neutrinos, Capri (1997); hep-ph/9710273.

- [12] Y. Fukuda *et al.*, Phys. Lett. **335B**, 237 (1994); **280B**, 146 (1992); **205B**, 416 (1988); D. Casper *et al.*, Phys. Rev. Lett. **66**, 2561 (1991); Phys. Rev. D **46**, 3720 (1992); W. Allison *et al.*, Phys. Lett. **391B**, 491 (1997).
- [13] C. Athanassopoulos *et al.*, Phys. Rev. C **54**, 2685 (1996); Phys. Rev. Lett. **77**, 3082 (1996); nucl-ex/9709006.